

Tracing the evolution of massive galaxies up to $z \sim 3$

M. Longhetti¹, P. Saracco¹, S. Cristiani², A. Fontana³, E. Giallongo³, M. Nonino², and E. Vanzella⁴

¹ INAF - Osservatorio Astronomico di Brera, Milano, Italy

² European Southern Observatory

³ INAF - Osservatorio Astronomico di Roma, Italy

⁴ INAF - Osservatorio Astronomico di Trieste, Italy

1 J-K > 3 galaxies: unveiling $M > 10^{11} M_{\odot}$ early-type galaxies at $z \sim 3$

Objects with unusual red near-IR color $J-K > 3$ are extremely rare at magnitude brighter than $K=20$, while their surface density increases at fainter magnitudes. The nature of these sources has not yet been firmly established even if all the analysis performed so far conclude that they are galaxies at $z > 2-3$ ([1], [2], [3]). A detailed analysis based on a multi-band data set (from $0.3\mu\text{m}$ to $2.15\mu\text{m}$) of three $J-K > 3$ sources selected at $K_s < 22$ on the HDF-S ([4]) has shown that: i) the three galaxies are at redshift $2.5 < z < 3$; ii) they have a stellar mass content of $M_{star} \sim 10^{11} M_{\odot}$; iii) they would populate the bright end ($L_{z=0} \sim L^*$) of the local luminosity function of galaxies even assuming they evolve passively; iv) they cannot follow an evolution significantly different from passive aging.

2 Depicting the evolution of massive galaxies from $z=3$ to $z=0$

Thus, these $J-K > 3$ objects selected are the $z \sim 3$ counterpart of local massive early-type galaxies. Between $z=3$ and $z=0$ we expect to find the same population of galaxies at a different evolutionary status, namely with older stellar content. For instance, at $z=1-1.5$ we expect to find early-type galaxies ($t_{z=3} - t_{z=1.5} \approx 2$ Gyr older. Such old early-type galaxies have been found among the EROs selected with $R-K_i > 5$ (e.g.[5]). In order to find their counterpart at the intermediate redshift $z \sim 2$, we have identified a suitable color selection criterion by exploring the the expected colors evolution as derived by synthetic models ([6]). The results are displayed in Fig. 1: for short SF time scale ($\tau < 1$ Gyr galaxies at $z \sim 2$ have colors $J-K < 3$ and $I-H > 4$). In fact, adopting these selection criteria on the HDF-S data, we identified 4 massive evolved galaxies at $1.5 < z < 2.5$ which seem to be the link between the $J-K > 3$ galaxies ($z \sim 3$) and the EROs ($z \sim 1$). An example of this passive evolution connection is proposed in the right panel of Fig. 1: 822 (upper panel) a $J-K > 3$ galaxy selected in the HDF-S well described by a young SSP at $z \sim 2.9$; 1203 (middle panel) one of the four massive ($M_{star} \geq 10^{11}$

$M_{\odot}$) galaxies selected in the HDF-S on the basis of the selection criteria above described, well described by a 1.4 Gyr old SSP at $z \sim 1.8$; S7F5.254 (lower panel) the spectrum of a massive early-type galaxy at $z=1.2$ ([5]) superimposed on the template of a 3 Gyr old SSP.

The photometry, the stellar masses and the mean ages of the stellar populations of these three massive galaxies at different redshift are apparently those we expect from a massive galaxy fully assembled at $z \sim 3$ which evolves passively in time. This result is based, in fact, on a couple of objects and may be simply a coincidence. However, it is worthy to note that the number density of the three classes of objects represented in the figure and that of the local massive galaxies are well compatible with the depicted scenario. Actually, the number density of $J-K > 3$ and $K < 22$ galaxies ($2.5 < z < 3$, $M_{star} \geq 10^{11} M_{\odot}$ galaxies) is $\rho = 1.2 \times 10^{-4} \text{ Mpc}^{-3}$ ([4]) and that of $J-K < 3$ and $I-H > 4$ ($1.5 < z < 2.5$) calculated from the HDF-S data results to be the same within the errors. The number density of the early-type galaxies at $1.0 < z < 1.5$ ($R-K > 5$) is consistent with the previous value once the same stellar mass threshold ($M_{star} \geq 10^{11} M_{\odot}$) is taken into account ([4]). The local density of $M_{star} \geq 10^{11} M_{\odot}$ elliptical galaxies is $\rho \simeq 3 \times 10^{-4} \text{ Mpc}^{-3}$ ([7]): at least 40% of the local population of bright massive early-type galaxies can be explained as the results of a simple passive evolution started from galaxies already assembled at $z \sim 3$.

References

1. M. Dickinson, C. Hanley, R. Elston, et al. 2000, ApJ, 531, 624
2. P.B. Hall, M. Sawicki, P. Martini, et al. 2001, ApJ, 121, 1840
3. T. Totani, Y. Yoshii, F. Iwamuro, et al. 2001, ApJ, 558, L87
4. P. Saracco, M. Longhetti, E. Giallongo, et al. 2003, astro-ph/0310131
5. P. Saracco, M. Longhetti, P. Severgnini, et al. 2003, A&A, 398, 127
6. M. Longhetti, P. Saracco, E. Giallongo, et al. 2004, in preparation
7. R.O. Marzke, L.N. Da Costa, P.S. Pellegrini, et al. 1998, ApJ, 503, 617




Fig. 1. *left panel:* each line represents the evolution with redshift of the J-K and I-H colors for different SFHs between $z=1$ (filled circles), $z=2$ (filled triangles) and $z=3$ (filled squares) (solar metallicity, Salpeter IMF, formation redshift $z_f=6$); *right panel:* possible passive evolution scenario (details in the text)